

# ChemComm

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: M. L. Odyniec, A. C. Sedgwick, A. Swan, M. Weber, T. M. S. Tang, J. E. Gardiner, M. Zhang, Y. Jiang, G. Kociok-Kohn, R. B. P. Elmes, S. Bull, X. He and T. D. James, *Chem. Commun.*, 2018, DOI: 10.1039/C8CC04316G.




This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [author guidelines](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the ethical guidelines, outlined in our [author and reviewer resource centre](#), still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

A novel fluorescence-based scaffold was developed enabling the production of 'AND' based fluorescence probes - ROS/RNS and a second analyte





Journal Name

COMMUNICATION

## 'AND' based fluorescence scaffold for the detection of ROS/RNS and a second analyte

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Maria L. Odyniec<sup>a†</sup>, Adam C. Sedgwick<sup>a\*†</sup>, Alexander H. Swan<sup>a</sup>, Maria Weber<sup>a</sup>, T. M. Simon Tang<sup>a</sup>, Jordan E. Gardiner<sup>a</sup>, Miao Zhang<sup>b</sup>, Yun-Bao Jiang<sup>b</sup>, Gabriele Kociok-Kohn<sup>a</sup>, Robert. B. P. Elmes<sup>c</sup>, Steven D. Bull<sup>a\*</sup>, Xiao-Peng He<sup>d\*</sup> and Tony D. James<sup>a\*</sup>

Biological processes often require more than one chemical species to achieve the desired outcome. Traditionally, fluorescence probes have focused on the detection of a single biomarker for a specific process. In this work, we set out to develop a number of fluorescence probes that enable the detection of a chosen analyte in the presence of reactive oxygen/nitrogen species (ROS/RNS). These fluorescence probes when activated result in the formation of the highly fluorescent pink dye, resorufin. Therefore, we have labelled these fluorescent probes as 'pinkments'. Our first 'Pinkment' was shown to detect biologically relevant concentrations of ONOO<sup>-</sup> and have an excellent selectivity against other ROS/RNS. Pinkment-OH was developed to provide a core unit which could be easily functionalised to produce a range of 'AND' based fluorescence probes for the detection of ROS/RNS and a second analyte. For proof of concept, we synthesised Pinkment-OTBS and Pinkment-OAc. These 'AND'-based probes were successfully shown to detect ROS/RNS and F<sup>-</sup> or esterase, respectively.

Since the discovery of the first fluorescence-based probe in 1867, fluorescence probes have revolutionised the understanding of biological systems.<sup>1-4</sup> Historically, these probes have focused on the detection of a single analyte or biomarker. However, biological systems are complex with more than one chemical species being released/present during any biological processes. For example, glutathione (GSH) accumulates at the nucleus during the cell cycle to aid transcription factors binding to DNA<sup>5</sup> and the pathological role

of Zn<sup>2+</sup> is believed to be associated with the glutamate system.<sup>6</sup> Furthermore, the sensitivity of a cell towards peroxynitrite (ONOO<sup>-</sup>) largely depends on the concentration of intracellular GSH.<sup>7-10</sup> Therefore, in order to further understand cellular functions and the root causes of disease it is important to be able to study biologically important species simultaneously.

Alongside the development of the field of fluorescence probes, the field of molecular logic gates has developed.<sup>11</sup> Molecular logic gates are molecules that have the ability to bind or react with multiple analytes (input) and turn it into a measurable optical output. Consequently, these attractive molecules are now emerging in the literature demonstrating the ability to simultaneously detect multiple analytes in biological systems.<sup>12-17,18-21</sup> Within our research group, we are interested in developing reaction-based fluorescence probes including 'AND'-based fluorescence probes for the detection of biologically important analytes.<sup>22-24</sup> Dual responsive ('AND') fluorescence probes require both analytes being present to produce a fluorescence response. In this work, we identified a previously reported boronate-based fluorescence probe developed by Chang *et al.* **PR1**, with a free amino group attached (Fig.1). Boronates have well-known reactivity towards ONOO<sup>-</sup> and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>),<sup>25</sup> therefore we believed **PR1** could provide a suitable scaffold for the development of 'AND'-based systems for the detection of ROS/RNS and an second analyte.<sup>26</sup> In this work, we initially functionalised **PR1** with an additional ROS/RNS trigger, benzyl boronic acid ester, which afforded a selective ONOO<sup>-</sup> probe known as **Pinkment** (Fig. 1).

<sup>a</sup> Department of Chemistry, University of Bath, Bath, BA2 7AY, UK. Email: t.d.james@bath.ac.uk; s.d.bull@bath.ac.uk

<sup>b</sup> Department of Chemistry, College of Chemistry and Chemical Engineering, and the MOE Key Laboratory of Analytical Sciences, Xiamen University, Xiamen, 361005, China Drug &

Department of Chemistry, Maynooth University, Maynooth, Co. Kildare, Ireland

<sup>c</sup> Key Laboratory for Advanced Materials & Feringa Nobel Prize Scientist Joint, Research Center, East China University of Science and Technology, 130 Meilong Rd., Shanghai 200237, P. R. China.


<sup>†</sup>ACS and MLO contributed equally, ACS - Organic Synthesis, MLO -Fluorescence analysis

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x



## COMMUNICATION


## Journal Name



**Fig. 1** – PR1 and **Pinkment** fluorescence probes for the detection of reactive oxygen and reactive nitrogen species (ROS/RNS). **Pinkment-Bn** demonstrates the requirement elimination to produce a free N-H.

The current literature route to obtain **PR1** involves a very low yielding first step (10 % overall yield 7 %), which uses harsh conditions (hydrobromic acid (HBr)).<sup>26</sup> Therefore, we alkylated commercially available phenoxazine using NaH and 4-bromobenzyl bromide to afford **1** in excellent yield (78 %). **1** was then subsequently dissolved in  $\text{CHCl}_3$  and NBS was added portion wise to afford dibrominated intermediate **2** in good yield (48 %), which required only trituration (EtOAc) for purification. Lastly, Suzuki-Miyaura conditions were applied to **2** using bis(pinacolato)diboron ( $\text{B}_2\text{pin}_2$ ) as a boron transfer agent to furnish **Pinkment** in 36 % yield (See ESI – Scheme S4). In the presence of ONOO<sup>-</sup>, all three boronates were oxidised on **Pinkment**, forming the highly fluorescent pink dye, resorufin (See ESI – Scheme S1) and a large fluorescence increase was observed at 590 nm (See ESI – Fig. S1). **Pinkment** was able to detect biologically relevant concentrations of ONOO<sup>-</sup><sup>27,28</sup> and displayed excellent selectivity towards ONOO<sup>-</sup> (See ESI – Fig. S2-4) To demonstrate the requirement of an ‘eliminating’ benzyl group on the free N-H, **Pinkment-Bn** (Fig. 1) was synthesised (See ESI – Scheme S5) and as expected no increase in fluorescence intensity was observed with the addition of ONOO<sup>-</sup> (10  $\mu\text{M}$ ) (See ESI – Fig. S5).

From these results, we believed we could develop a range of ‘AND’-based **Pinkment** fluorescence probes by altering the functionality on the benzyl group to react with a specific target analyte. To achieve this, we developed a synthesis for **Pinkment-OH** (Fig.2), which provides a novel fluorescence core probe that can be functionalised with any chosen reactive chemical trigger for the detection of a chosen analyte in the presence of ROS/RNS (Fig. 2a).




**Fig. 2** – (a) **Pinkment-OH** – a core fluorescence unit that enables the synthesis of ‘AND’ based fluorescence probes for the detection of (ROS/RNS) and a second analyte. (b) Molecular structure of **Pinkment-OH** with thermal ellipsoids shown at 50% probability. Hydrogen atoms are omitted for clarity except for the OH hydrogen atom. (c) **Pinkment-OTBS** and **Pinkment-OAc** for the detection of ROS/RNS ‘AND’ fluoride or esterase, respectively. see ESI-4). **CCDC 1844385** contain the supplementary crystallographic data for **Pinkment-OH**. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).

For the synthesis of **Pinkment-OH**, the phenol of 4-hydroxybenzyl alcohol was selectively *piv* protected to afford **5** using pivaloyl chloride and  $\text{NEt}_3$  in DCM. **5** was then converted to its corresponding bromide **6** using  $\text{MsCl}$  and  $\text{NEt}_3$  followed by the addition of LiBr. Phenoxazine was then alkylated with **6** to produce **7** using NaH and DMF in satisfactory yield (51 %). **7** was then dibrominated using NBS in  $\text{CHCl}_3$  to afford **8**, which was subsequently deprotected using DIBAL-H to afford **9** in good yield (67 %). The same Suzuki-Miyaura conditions were then applied to furnish **Pinkment-OH** in excellent yield (83 %) (See ESI – Scheme S6). To illustrate the stability of **Pinkment-OH**, an x-ray structure was obtained (Fig. 2b). Interestingly, **Pinkment-OH** formed dimers via intermolecular hydrogen bonding (see ESI – Fig. S20).

**Pinkment-OH** was shown to have a high sensitivity and excellent selectivity towards ONOO<sup>-</sup> making it suitable for cellular imaging experiments (See ESI – Fig. S6-9). As proof of concept for ‘AND’-based pinkment sensors, we chose to synthesise **Pinkment-OTBS** and **Pinkment-OAc** (Fig. 2c and 2d), which can be used to detect ROS/RNS and fluoride ( $\text{F}^-$ ) or esterase, respectively.


To afford **Pinkment-OTBS**, **Pinkment-OH** was silyl protected using standard silyl protection conditions, however, the reaction was found to be very slow (3 d) and **Pinkment-OTBS** was isolated in very low yield (11 %). We then evaluated the ability of **Pinkment-OTBS** to detect  $\text{F}^-$  ‘AND’ ONOO<sup>-</sup> in a buffer solution (52% w/w MeOH:H<sub>2</sub>O). This MeOH/H<sub>2</sub>O buffer solution was required for the silyl-ether deprotection reaction to proceed effectively. As shown in ESI – Fig. S10, 100 % PBS produced a much smaller response when both analytes were added, which is consistent with literature reported TBS fluorescence probes.<sup>29,30</sup> As shown in Fig. 3, addition of tert-butylammonium fluoride (TBAF - Source of  $\text{F}^-$ ) led to no increase in fluorescence intensity, however, subsequent additions of ONOO<sup>-</sup> (0 – 20  $\mu\text{M}$ ) led to a significant increase in fluorescence intensity.





**Fig. 3** – Fluorescence spectra of **Pinkment-OTBS** (5  $\mu\text{M}$ ) with addition of TBAF (10 mM) and incubated for 1 h and then measured. Followed by incremental additions of  $\text{ONOO}^-$  (0 – 20  $\mu\text{M}$ ) The data was obtained in PBS (52% w/w MeOH:H<sub>2</sub>O), pH = 8.2 at 25  $^\circ\text{C}$ .  $\lambda_{\text{ex}}$ =545-20 nm.


To ensure both analytes were required, the addition of TBAF and  $\text{ONOO}^-$  were then repeated in reverse order. Due to the high reactivity of  $\text{ONOO}^-$ ,<sup>8, 31</sup> a small increase in fluorescence intensity was observed on addition to **Pinkment-OTBS** (20  $\mu\text{M}$ ). However, like Fig.3, a large increase in fluorescence intensity was only observed after the subsequent addition of TBAF (0 – 10 mM). As shown in ESI - Fig. S12 and S13, **Pinkment-OTBS** displayed an excellent selectivity towards  $\text{ONOO}^-$  against other ROS in the presence of TBAF. **Pinkment-OTBS** was then screened against other halide sources (TBAB, TBAC and TBAI) in the presence of  $\text{ONOO}^-$  (20  $\mu\text{M}$ ) see ESI-Fig. S14. As expected **Pinkment-OTBS** displayed excellent selectivity towards TBAF in the presence of  $\text{ONOO}^-$ .



**Fig. 4** - Fluorescence spectra of **Pinkment-OTBS** (5  $\mu\text{M}$ ) with the addition of  $\text{ONOO}^-$  (20  $\mu\text{M}$ ) followed by incremental additions of TBAF (0 – 10 mM) measurements were taken after 90 mins of both additions. The data was obtained in PBS (52% w/w MeOH:H<sub>2</sub>O), pH = 8.2 at 25  $^\circ\text{C}$ .  $\lambda_{\text{ex}}$ =545-20 nm. Dashed line represents  $\text{ONOO}^-$  (20  $\mu\text{M}$ ) addition only


We then turned our attention towards the synthesis of **Pinkment-OAc**. Despite the poor reactivity with TBS-Cl, **Pinkment-OH** reacted with AcCl smoothly and produced **Pinkment-OAc** in excellent yield (71 %). Due to the known ability of  $\text{ONOO}^-$  to react with carbonyls, its addition to **Pinkment-OAc** was avoided (See ESI – S15 and S16). Therefore,

as proof of concept, we evaluated **Pinkment-OAc** with H<sub>2</sub>O<sub>2</sub> 'AND' F. As shown in Fig. 5, the addition of porcine liver esterase, PLE (0.6 U) led to no increase in fluorescence intensity. However, subsequent additions of H<sub>2</sub>O<sub>2</sub> (0 – 1 mM) led to a large increase in fluorescence.



**Fig. 5** - Fluorescence spectra of **Pinkment-OAc** (5  $\mu\text{M}$ ) for the incubation (15 min) of porcine liver esterase, PLE (0.6 U) followed by incremental additions of H<sub>2</sub>O<sub>2</sub> (0 – 1 mM) measurements were taken after 1 h. The data was collected in PBS (100% w/w H<sub>2</sub>O) at pH=7.4 ,25  $^\circ\text{C}$ .  $\lambda_{\text{ex}}$ =545-20 nm.

Again, to ensure that both analytes were required, the concentration of H<sub>2</sub>O<sub>2</sub> was kept constant (1 mM) and the amount of PLE was varied. Due to the known ability of H<sub>2</sub>O<sub>2</sub> to hydrolyse carbonyl containing compounds, it was no surprise that there was a small increase in fluorescence intensity.<sup>32, 33</sup> However, with increasing additions of PLE much larger increases in fluorescence intensity were observed (Fig. 6). Interestingly, concentrations > 0.6 U led to a smaller increase in fluorescence. This is believed to be the result of the excess esterase reacting with the added H<sub>2</sub>O<sub>2</sub>.



**Fig. 6** - Fluorescence spectra of **Pinkment-OAc** (0.5  $\mu\text{M}$ ) following incubation (15 mins) with PLE (0.1 - 0.6 U) and subsequent addition of H<sub>2</sub>O<sub>2</sub> (1 mM). The dotted line represents the response to hydrogen peroxide on its own. The data was collected in PBS buffer (100% w/w H<sub>2</sub>O) at pH=7.4 ,25  $^\circ\text{C}$ .  $\lambda_{\text{ex}}$ =545-20 nm Dashed line represents H<sub>2</sub>O<sub>2</sub> (1 mM) addition only

In summary, several fluorescence-based probes have been synthesised that when activated result in the formation of the highly fluorescent dye, resorufin. As the dye is pink in colour, we have labelled these fluorescent probes as 'pinkments'. The





original **Pinkment** was shown to detect biologically relevant concentrations of ONOO<sup>-</sup> and have an excellent selectivity against other ROS/RNS making it suitable for cellular imaging experiments. **Pinkment** provided a suitable platform for the development of 'AND'-based fluorescent probes for the detection of ROS/RNS and another analyte. **Pinkment-OH** is a core building block that enables easy functionalisation in order to produce a range of 'AND' based fluorescence probes for the detection of (ROS/RNS) and a second analyte. As proof of concept, we developed **Pinkment-OTBS** and **Pinkment-OAc** for the detection of ROS/RNS and F<sup>-</sup> and esterase, respectively. Both **Pinkment-OTBS** and **Pinkment-OAc** were demonstrated to be successful 'AND'-based fluorescence probes. However, both probes demonstrated slight reactivity to ROS/RNS alone, due to both silyl ether and acetate functional groups being unstable towards nucleophiles. We are now turning our attention to the attachment of palladium ion (Pd<sup>2+</sup>) and mercury ion (Hg<sup>2+</sup>) reactive triggers, previously developed by Ahn *et al.* and Koide *et al.*<sup>34, 35</sup> It is believed that the alkyl ethers will provide stability towards ROS/RNS in addition it is known that Hg<sup>2+</sup> and Pd nanoparticles produce ROS in biological systems promoting apoptosis.<sup>36-39</sup>

### Acknowledgements

We would like to thank the EPSRC and the University of Bath for funding for ACS, MLO, JEG for studentships. MW would like to thank the EPSRC for funding (i) EP/L016354/1 and CDT in Sustainable Chemical Technologies. TDJ wishes to thank the Royal Society for a Wolfson Research Merit Award. M Zhang and Y-B Jiang were supported by the NSF of China under grants 21435005 and 21521004. X.-P. He wishes to thank the National Natural Science Foundation of China (21722801), the Science and Technology Commission of Shanghai Municipality (15540723800) and the Shanghai Rising-Star Program (16QA1401400) for financial support. NMR characterisation facilities were provided through the Chemical Characterisation and Analysis Facility (CCAF) at the University of Bath ([www.bath.ac.uk/ccaf](http://www.bath.ac.uk/ccaf)). The EPSRC UK National Mass Spectrometry Facility at Swansea University is thanked for analyses. All data supporting this study are provided as supplementary information accompanying this paper (ESI†).

### Conflicts of interest

No conflicts of interest

### Notes and references

1. D. Wu, A. C. Sedgwick, T. Gunnlaugsson, E. U. Akkaya, J. Yoon and T. D. James, *Chem. Soc. Rev.*, 2017, **46**, 7105-7123.
2. J. Chan, S. C. Dodani and C. J. Chang, *Nat. Chem.*, 2012, **4**, 973-984.
3. Y. M. Yang, Q. Zhao, W. Feng and F. Y. Li, *Chem. Rev.*, 2013, **113**, 192-270.

4. X. P. He and H. Tian, *Chem*, 2018, **4**, 246-268.
5. J. Markovic, C. Borrás, A. Ortega, J. Sastre, J. Vina and F. V. Pallardo, *J. Biol. Chem.*, 2007, **282**, 20416-20424.
6. B. Pochwat, G. Nowak and B. Szewczyk, *Neural Plast.*, 2015, DOI: 10.1155/2015/591563.
7. K. A. Marshall, R. Reist, P. Jenner and B. Halliwell, *Free Radic. Biol. Med.*, 1999, **27**, 515-520.
8. P. Pacher, J. S. Beckman and L. Liaudet, *Physiol. Rev.*, 2007, **87**, 315-424.
9. J. P. Bolanos, S. J. R. Heales, J. M. Land and J. B. Clark, *J. Neurochem.*, 1995, **64**, 1965-1972.
10. M. Nakamura, V. H. Hourani, R. S. Ronson, D. A. Velez, X. L. Ma, S. Katzmark, J. Robinson, L. S. Schmarkey, Z. Q. Zhao, N. P. Wang, R. A. Guyton and J. Vinten-Johansen, *Circulation*, 2000, **102**, 332-338.
11. S. Erbas-Cakmak, S. Kolemen, A. C. Sedgwick, T. Gunnlaugsson, T. D. James, J. Yoon and E. U. Akkaya, *Chem. Soc. Rev.*, 2018, **47**, 2228-2248.
12. C. Y. Ang, S. Y. Tan, S. J. Wu, Q. Y. Qu, M. F. E. Wong, Z. Luo, P. Z. Li, S. T. Selvan and Y. L. Zhao, *J. Mater. Chem. C*, 2016, **4**, 2761-2774.
13. S. Debieu and A. Romieu, *Org. Biomol. Chem.*, 2015, **13**, 10348-10361.
14. G. C. Van de Bittner, C. R. Bertozzi and C. J. Chang, *J. Am. Chem. Soc.*, 2013, **135**, 1783-1795.
15. F. B. Yu, P. Li, B. S. Wang and K. L. Han, *J. Am. Chem. Soc.*, 2013, **135**, 7674-7680.
16. X. F. Yang, Q. Huang, Y. G. Zhong, Z. Li, H. Li, M. Lowry, J. O. Escobedo and R. M. Strongin, *Chem. Sci.*, 2014, **5**, 2177-2183.
17. S. Resa, A. Orte, D. Miguel, J. M. Paredes, V. Puente-Munoz, R. Salto, M. D. Giron, M. J. Ruedas-Rama, J. M. Cuerva, J. M. Alvarez-Pez and L. Crovetto, *Chem. Eur. J.*, 2015, **21**, 14772-14779.
18. X. P. He, X. L. Hu, T. D. James, J. Yoon and H. Tian, *Chem. Soc. Rev.*, 2017, **46**, 6687-6696.
19. L. Yu, S. L. Wang, K. Z. Huang, Z. G. Liu, F. Gao and W. B. Zeng, *Tetrahedron*, 2015, **71**, 4679-4706.
20. J. L. Kolanowski, F. Liu and E. J. New, *Chem. Soc. Rev.*, 2018, **47**, 195-208.
21. A. Romieu, *Org. Biomol. Chem.*, 2015, **13**, 1294-1306.
22. A. C. Sedgwick, H. H. Han, J. E. Gardiner, S. D. Bull, X. P. He and T. D. James, *Chem. Commun.*, 2017, **53**, 12822-12825.
23. A. C. Sedgwick, R. S. L. Chapman, J. E. Gardiner, L. R. Peacock, G. Kim, J. Yoon, S. D. Bull and T. D. James, *Chem. Commun.*, 2017, **53**, 10441-10443.
24. A. C. Sedgwick, H. H. Han, J. E. Gardiner, S. D. Bull, X. P. He and T. D. James, *Chem. Sci.*, 2018, **9**, 3672-3676.
25. A. Sikora, J. Zielonka, M. Lopez, J. Joseph and B. Kalyanaraman, *Free Radic. Biol. Med.*, 2009, **47**, 1401-1407.
26. E. W. Miller, A. E. Albers, A. Pralle, E. Y. Isacoff and C. J. Chang, *J. Am. Chem. Soc.*, 2005, **127**, 16652-16659.
27. X. M. Chen, B. H. Zhou, T. T. Yan, H. Wu, J. H. Feng, H. S. Chen, C. Gao, T. Peng, D. Yang and J. G. Shen, *Free Radic. Biol. Med.*, 2018, **117**, 158-167.
28. H. Fujigaki, K. Saito, F. Lin, S. Fujigaki, K. Takahashi, B. M. Martin, C. Y. Chen, J. Masuda, J. Kowalak, O. Takikawa, M. Seishima and S. P. Markey, *J. Immunol.*, 2006, **176**, 372-379.
29. X. F. Yang, S. J. Ye, Q. Bai and X. Q. Wang, *J. Fluoresc.*, 2007, **17**, 81-87.
30. J. Cao, C. C. Zhao, P. Feng, Y. L. Zhang and W. H. Zhu, *RSC Adv.*, 2012, **2**, 418-420.
31. D. Yang, Y. C. Tang, J. Chen, X. C. Wang, M. D. Bartberger, K. N. Houk and L. Olson, *J. Am. Chem. Soc.*, 1999, **121**, 11976-11983.
32. M. Abo, Y. Urano, K. Hanaoka, T. Terai, T. Komatsu and T. Nagano, *J. Am. Chem. Soc.*, 2011, **133**, 10629-10637.
33. J. Yang, X. L. Zhang, P. Yuan, Y. G. Xu, J. Grutzendler, Y. H. Shao, A. Moore and C. Z. Ran, *Proc. Natl. Acad. Sci. U. S. A.*, 2017, **114**, 12384-12389.
34. M. Santra, S. K. Ko, I. Shin and K. H. Ahn, *Chem. Commun.*, 2010, **46**, 3964-3966.
35. F. L. Song, S. Watanabe, P. E. Floreancig and K. Koide, *J. Am. Chem. Soc.*, 2008, **130**, 16460-16461.
36. B. J. Shenker, T. L. Guo and I. M. Shapiro, *Environ. Res.*, 2000, **84**, 89-99.
37. A. E. A. Moneim, *Neural Regen. Res.*, 2015, **10**, 881-882.
38. A. E. A. Moneim, *Metab. Brain Dis.*, 2015, **30**, 935-942.
39. S. Alarifi, D. Ali, S. Alkahtani and R. S. Almeer, *Oxid. Med. Cell. Longev.*, 2017, DOI: 10.1155/2017/8439098.

